

Possible CPT Violation from Planck Scale Effects

Bipin Singh Koranga^{a*}, Mohan Narayan^{b†} and S. Uma Sankar^{a‡}

^a*Department of Physics Indian Institute of Technology, Bombay, Mumbai 400 076*

^b*Department of Physics, Mumbai University*

Institute of Chemical Technology, Mumbai 400 019

Abstract

At present there is good agreement between the neutrino mass-squared difference determined from the solar neutrino data and the anti-neutrino mass-squared difference determined from the KamLAND reactor anti-neutrino experiment. However, the central values of the two cases differ from each other by about 10^{-5} eV^2 . An improvement in the accuracy of both the solar neutrino experiments and reactor anti-neutrino experiments can establish the existence of a non-zero difference between neutrino and anti-neutrino mass-squared differences and provide a signal for CPT violation. In this paper, we show how such a difference can arise through the CPT violating neutrino mass terms from Planck scale physics.

* bipinkar@phy.iitb.ac.in

† mohan@udct.org

‡ uma@phy.iitb.ac.in

I. INTRODUCTION

KamLAND experiment recently detected the distortion due to oscillations in the anti-neutrino spectrum from the reactors and determined the corresponding mass-square difference, $\overline{\Delta}_{21}$, to great precision [1]. At present there is good agreement between $\overline{\Delta}_{21}$ and the mass-squared difference of the neutrinos, Δ_{21} , determined from the analysis of solar neutrino data [2]. However, the best-fit values of the two Δ s differ from each other by 10^{-5} eV². Together, solar and KamLAND data impose the constraint $|\Delta_{21} - \overline{\Delta}_{21}| \leq 1.1 \times 10^{-4}$ eV² [3]. Future reactor experiments, located at a distance of about 70 Km from the source so that the oscillation minimum coincides with spectral maximum, are expected to improve the precision of $\overline{\Delta}_{21}$ even further [4]. Similarly future solar neutrino experiments, such as LENS [5] and other experiments [6], are expected to improve the accuracy of Δ_{21} . These future experiments may indeed show that there is a non-zero difference between Δ_{21} and $\overline{\Delta}_{21}$, thus establishing a signal for CPT violation in the neutrino sector [7, 8].

If Δ_{21} and $\overline{\Delta}_{21}$ are indeed found to be different, a natural question to ask is: How does this CPT violation arise? In this letter, we assume that CPT violation in neutrino sector arises due to Planck scale effects. We parametrize these effects in terms of Planck scale CPT violating neutrino mass terms and calculate the difference between Δ_{21} and $\overline{\Delta}_{21}$ arising due to these terms.

We assume that neutrino masses mainly arise due to Grand Unified Theory (GUT) dynamics via see-saw mechanism [9] and these masses are CPT conserving. We further assume that the CPT violation arises only at Planck scale and parametrize it by the effective neutrino mass term

$$\mathcal{M}_{\text{CPT}} = \frac{v^2}{2M_{\text{Pl}}} \lambda_{\alpha\beta} \quad (1)$$

where α and β are flavour indices. The mass term for anti-neutrinos will have the opposite sign [10]. Since these are effective masses arising from Planck scale effects, they are suppressed by $1/M_{\text{Pl}}$. Since these are assumed to be the masses at the low energy scale, the electroweak vacuum expectation value (VEV) $v = 174$ GeV is used to make these terms have dimension of mass. In eq. (1) the term λ is a 3×3 matrix in flavour space whose elements are of order 1. We assume that the Planck scale interaction is “flavour blind”, *i.e.* the elements of the matrix λ are independent of α, β indices. In this case, the contribution to

the neutrino mass matrix is of the form:

$$\mu \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \quad (2)$$

where the scale μ is

$$\mu = v^2/2M_{\text{Pl}} \simeq 10^{-6} \text{ eV}. \quad (3)$$

In our calculations, we take eq. (2) as a perturbation to the main part of the neutrino mass matrix, that is generated by GUT dynamics.

II. CALCULATION

The theoretical framework, in which the Planck scale mass terms are treated as perturbation to GUT scale neutrino masses, is developed in ref. [11]. Here we briefly recapitulate some salient features of this framework. In the discussion below the labels α and β refer to flavour eigenstates and the labels i and j refer to mass eigenstates. The unperturbed (0^{th} -order) neutrino mass matrix $\mathcal{M}$ is diagonalized by a unitary matrix U to yield the matrix M , whose eigenvalues are real and non-negative. As stated before, $\mathcal{M}$ is generated by grand unified dynamics and is related to U and M through the relation

$$\mathcal{M} = U^* M U^\dagger, \text{ where } U = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu1} & U_{\mu2} & U_{\mu3} \\ U_{\tau1} & U_{\tau2} & U_{\tau3} \end{pmatrix}. \quad (4)$$

We adopt the usual parameterization [12]: $U_{e2}/U_{e1} = \tan \theta_{12}$, $U_{\mu3}/U_{\tau3} = \tan \theta_{23}$ and $|U_{e3}| = \sin \theta_{13}$. We include all possible phases in the definition of the neutrino mixing matrix,

$$U = \text{diag}(e^{if_i}) R(\theta_{23}) \Pi R(\theta_{13}) \Pi^* R(\theta_{12}) \text{diag}(e^{ia_i}). \quad (5)$$

The phase δ appearing in $\Pi = \text{diag}(e^{i\delta/2}, 1, e^{-i\delta/2})$ is the one that enters oscillation probabilities and leads to CP violation in neutrino oscillations. a_i are the so called Majorana phases and the matrix with this phases has the form $\text{diag}(e^{ia_1}, e^{ia_2}, 1)$. The diagonal phase matrix on the left is given by $\text{diag}(e^{if_1}, e^{if_2}, e^{if_3})$, which are usually absorbed into the definition of the respective charged lepton field. It is possible to rotate away the phases f_i , if the mass

matrix (4) *is the complete mass matrix*. However, since we are going to add another contribution to this mass matrix, the phases f_i of the zeroth order mass matrix have an impact on the complete mass matrix and thus must be retained. By the same token, the Majorana phases which are usually redundant for oscillations have a dynamical role to play now.

Planck scale effects will add other contributions to the mass matrix. The additional term has the form given in eq. (2) and with its inclusion the neutrino mass matrix becomes

$$\mathcal{M} \rightarrow \mathcal{M}' = \mathcal{M} + \mu \lambda, \quad (6)$$

It is possible that the CPT violating mass terms may be arise not at Planck scale but at some scale M_X below M_{Pl} but well above the GUT scale (by an order of magnitude or so). In such a case, the perturbation parameter $\mu = v^2/2M_X$, rather than $v^2/2M_{pl}$.

We now define the hermitian matrix $\mathcal{M}'^\dagger \mathcal{M}'$ and find its eigenvalues and eigenvectors. The differences of pairs eigenvalues give us the modified mass-squared differences and the eigenvectors give us the modified neutrino mixing matrix. To first order in the small parameter μ , we have

$$\mathcal{M}'^\dagger \mathcal{M}' = \mathcal{M}^\dagger \mathcal{M} + \mu \lambda^\dagger \mathcal{M} + \mathcal{M}^\dagger \mu \lambda. \quad (7)$$

This matrix is diagonalized by a new unitary mixing matrix U' . We denote this diagonal matrix, correct to first order in μ , to be M'^2 . Using eq. (4), we can rewrite $\mathcal{M}$ in the above equation in terms of the diagonal matrix M . Converting $\mathcal{M}'$ also into its diagonal form M' , we can rewrite the above equation as

$$U' M'^2 U'^\dagger = U (M^2 + m^\dagger M + M m) U^\dagger \text{ with } m = \mu U^t \lambda U. \quad (8)$$

From the above equation, it follows that the mixing matrix U' , correct to first order in μ , is related to zeroth order mixing matrix U by

$$U' = U (1 + i\delta\theta), \quad (9)$$

where $\delta\theta$ is a hermitian matrix proportional to μ .

Substituting the expression for U' from eq. (9) in eq. (8) we obtain

$$M^2 + m^\dagger M + M m = M'^2 + [i\delta\theta, M'^2]. \quad (10)$$

Because M'^2 is diagonal, the diagonal terms of $[i\delta\theta, M'^2]$ in the above equation are zero. Therefore, to first order in μ , the mass squared differences $\Delta M_{ij}^2 = M_i^2 - M_j^2$ get modified

as:

$$\Delta M'_{ij}{}^2 = \Delta M_{ij}^2 + 2 (M_i \text{Re}[m_{ii}] - M_j \text{Re}[m_{jj}]), \quad (11)$$

where there is no summation over the repeated indices. The above equation shows the correction for neutrino mass-squared difference. Because the Planck scale corrections are assumed to be CPT violating, the correction for anti-neutrino mass-squared difference will have the opposite sign.

III. RESULTS

Note from eq.(11) that the correction term depends crucially on the type of neutrino mass spectrum. For a hierarchical or inverse hierarchical spectrum the correction is negligible. Hence we consider a degenerate neutrino spectrum and take the common neutrino mass to be 2 eV, which is the upper limit from the tritium beta decay experiment [13].

From the definition of the matrix m in eq. (8), we find

$$\begin{aligned} m_{11} &= \mu e^{i2a_1} (U_{e1} e^{if_1} + U_{\mu 1} e^{if_2} + U_{\tau 1} e^{if_3})^2 \\ m_{22} &= \mu e^{i2a_2} (U_{e2} e^{if_1} + U_{\mu 2} e^{if_2} + U_{\tau 2} e^{if_3})^2. \end{aligned} \quad (12)$$

The contribution of the terms in the Planck scale correction, $\text{Re}(m_{22}) - \text{Re}(m_{11})$, can be additive or subtractive depending on the values of the phases a_1 and a_2 . Similarly the magnitudes of $\text{Re}(m_{22})$ and $\text{Re}(m_{11})$ are functions of the phases f_i . Thus we try to find a combination of these phases which can give rise to a significant difference between neutrino and anti-neutrino mass-squared difference. In our calculations, we used $\theta_{12} = 34^\circ$, $\theta_{13} = 10^\circ$, $\theta_{23} = 45^\circ$ and $\delta_{cp} = 0$. We define the percentage correction P to be ratio of the difference between the corrected neutrino and anti-neutrino mass-squared difference to the unperturbed mass squared difference among the first and the second mass eigenstates as;

$$P = 100 \times (\Delta M'_{21}{}^2 - \overline{\Delta M'}_{21}{}^2) / \Delta M_{21}^2 \quad (13)$$

In Fig.(1) we plot our results as contours of constant P in the $a_1 - a_2$ plane, for $f_i = 0$. We note that the maximum difference allowed is about $\pm 5\%$ only. This is because the magnitudes of m_{22} and m_{11} remain relatively small in the limit $f_i = 0$. In Figs.(2) we plot contours of constant P in the $f_1 - f_2$ plane with $f_3 = 0 = a_1 = a_2$. When one phase is large and the other is small, there is a possibility that all the terms in m_{22} and m_{11} add up to

give a large value for their difference and hence a large value for P . We note that a much large change, varying from 30% to -20% is possible. In Figs.(3) and (4) we plot contours of constant P in $f_1 - f_3$ plane and in $f_2 - f_3$ plane. In generating these plots, we set all other phases to zero, as in the case of Fig.(2).

The effect scales directly as the common neutrino mass, so if one takes the WMAP [14] bound on the common neutrino mass which is 0.23 eV then there is negligible effect. However if there are new flavor blind interactions between GUT and Planck scales, then a low common mass is compensated by a lower scale of these new interactions and we still can get appreciable effects. In fact one can even say that if the gap between Δ_{21} and $\bar{\Delta}_{21}$ persists and future data indicate small neutrino mass, this can be an indication of new CPT violating flavor blind interactions below Planck scale.




FIG. 1: P as a function of the Majorana phases a_1 and a_2

IV. CONCLUSIONS

Both solar and reactor data are well accounted for by invoking neutrino oscillations. Fit to solar data give a large region for the neutrino mass squared difference in the two flavor




FIG. 2: P as a function of the phases f_1 and f_2

parameter space. The fit to reactor data however gives a very strongly constrained anti-neutrino mass squared difference. The best fits of the two mass squared differences are appreciably different from each other. Further improvement in KamLAND systematics and future solar neutrino data may remove this discrepancy. However if this mismatch between the best fits persists, then CPT breaking in the neutrino sector will be established. We have demonstrated that a flavour blind CPT violating neutrino masses from Planck scale physics can nicely accomodate this effect. This effect is crucially dependent on the neutrino mass spectrum and gives rise to observable difference between Δ_{21} and $\overline{\Delta}_{21}$ only for a degenerate neutrino mass spectrum with $m_\nu \simeq 2$ eV, which is the largest allowed value from tritium beta decay data. The low value of the common mass implied by the WMAP bound leads to negligible difference between Δ_{21} and $\overline{\Delta}_{21}$. This can however be compensated for by considering a slightly lower scale for the flavour blind CPT violating mass terms rather than the usual Planck scale.

Acknowledgement: We would like to thank Francesco Vissani for a critical reading of the




FIG. 3: P as a function of the phases f_1 and f_3

manuscript.

-
- [1] KamLAND Collaboration: T. Araki *et al*, Phys. Rev. Lett. **94**, 081801 (2005).
 - [2] J. N. Bahcall, M.C. Gonzalez-Garcia and C. Pena-Garay, JHEP **0408**:016 (2004).
 - [3] A. de Gouvea and C. Pena-Garay, Phys. Rev. D**71**, 093002 (2005);
A. de Gouvea, *Talk presented at Neutrino 2004, Paris*, hep-ph/0408246.
 - [4] S. Choubey and S. T. Petcov, Phys. Lett. **B 594**, 333 (2004);
H. Minakata, H. Nunokawa, W. J. C. Teves and R. Zukanovich Funchal, Phys. Rev. D**71**, 013005 (2005);
A. Bandyopadhyay, S. Choubey, S. Goswami and S. T. Petcov, Phys. Rev. D **72**, 033013 (2005).
S. Goswami, *Talk presented at LP 2005, Uppsala*, Int. J. of Mod. Phys. **A 21**, 1901 (2006). S. T. Petcov and T. Schwetz, hep-ph/0607155.




FIG. 4: P as a function of the phases f_2 and f_3

[5] T. Lasserre (for LENS Collaboration), Prog. Part. Nucl. Phys. **48**, 231 (2002).

[6] B. Vogelaar, Talk presented at Neutrino 2006, Santa Fe, June 2006.

[7] O. W. Greenberg, Phys. Rev. Lett. **89**, 231602 (2002).

[8] V. A. Kostelecky and M. Mewes, Phys. Rev. D **69** 016005 (2004).

[9] P. Minkowski, Phys. Lett. **B 67**, 421 (1977);

T. Yanagida, in *Proceedings of the workshop on unified theory and baryon number in the universe*, KEK, March 1979, eds. O. Sawada and A. Sugamoto;

M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, Stonybrook, 1979, eds. D. Freedman and P. van Nieuwenhuizen;

R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).

[10] G. Barenboim, L. Borissov, J. Lykken and A. Yu. Smirnov, JHEP **0210**, 001 (2002).

G. Barenboim, J. F. Beacom, L. Borissov and B. Kayser, Phys. Lett. **B537**, 227 (2002);

G. Barenboim and J. Lykken, Phys. Lett. **B554**, 73 (2003);

- [11] F. Vissani, M. Narayan and V. Berezhinsky, Phys. Lett. **B 571**, 209 (2003).
- [12] Review of Particle Physics, Phys. Lett. **B 592**, 145 (2004).
- [13] Ch. Weinheimer *et al*, Phys. Lett. **B 460**, 219 (1999);
V. M. Lobashev *et al*, Phys. Lett. **B 460**, 227 (1999).
- [14] M. Tegmark, Phys. Scripta **T 121**, 153 (2005); hep-ph/0503257